arXiv:1002.0445v3 [math.DG] 15 Aug 2010

A note about invariant Hermitian structures on adjoint orbits

Dmitri V. Alekseevsky and Liana David

August 17, 2010

Abstract: Let (g, J) be an invariant Hermitian structure on an adjoint orbit of a compact, connected, semisimple Lie group. We prove that if the Kähler form ω of (g, J) satisfies $dd^J\omega = 0$, then the pair (g, J) is Kähler. We apply this result to generalized Kähler geometry.

Key words: invariant Hermitian and Kähler structures, generalized Kähler structures, adjoint orbits.

2010 Mathematics Subject Classification: 53D18.

1 Introduction

A Hermitian structure on a smooth manifold M is a pair (g, J) where g is a Riemannian metric and J a g-skew-symmetric complex structure on M. The Hermitian structure (g, J) is Käher if the Kähler form $\omega(X, Y) := g(JX, Y)$ is closed. The main result in this note is the following:

Theorem 1. *Let (g, J) be an invariant Hermitian structure on an adjoint orbit of a compact connected semisimple Lie group. Let ω be the Kähler form of (g, J) and assume that*

$$dd^J\omega = 0. \tag{1}$$

Then the pair (g, J) is Kähler.

One motivation to consider condition (1) on Hermitian structures comes from generalized complex geometry, introduced by Nigel Hitchin in [6]. A generalized Kähler structure on a manifold M is usually defined as a pair $(\mathcal{J}_1, \mathcal{J}_2)$ of commuting generalized complex structures such that the composition $D := -\mathcal{J}_1\mathcal{J}_2$ satisfies an additional positivity condition (for the precise definition, see Section 4). According to [4], a generalized Kähler structure can also be defined as a bi-Hermitian structure (g, J_+, J_-) (i.e. (g, J_+) and (g, J_-) are Hermitian structures), together with a 2-form $b \in \Omega^2(M)$, usually called in the physical literature the b-field, such that

$$db = -d^{J_+}\omega_+ = d^{J_-}\omega_-, \tag{2}$$

where $\omega_\pm$ are Kähler forms of $(g, J_\pm)$. As an application of Theorem 1 we state:

Corollary 2. *Let (g, J_+, J_-, b) be an invariant generalized Kähler structure on an adjoint orbit of a compact, connected, semisimple Lie group. Then the pairs (g, J_+) and (g, J_-) are Kähler and $b \in \Omega^2(M)$ is closed.*

This note is organized as follows. In Section 2 we fix notations and we recall the description of invariant Hermitian and Kähler structures on adjoint orbits of compact semisimple Lie groups [1], [3], [8]. With these preliminaries, in Section 3 we prove Theorem 1. Finally, in Section 4 – intended for completeness of our exposition – we collect basic definitions from generalized complex geometry and we recall in more detail the relation mentioned above between generalized Kähler geometry and bi-Hermitian geometry [4].

Acknowledgements: We thank Paul Gauduchon for pertinent comments and suggestions on a first version of this paper and Radu Pantilie for his interest in this work. L. D. acknowledges financial support from a CNCSIS grant IDEI "Structuri geometrice pe varietati diferentiabile", code 1187/2008.

2 Preliminary material

Let G be a compact, connected, semisimple Lie group, with Lie algebra $\mathfrak{g}$, and $M = \{\mathrm{Ad}_g(X_0), g \in G\}$ an adjoint orbit of G, where $X_0 \in \mathfrak{g}$. We identify M with the coset space G/U, where $U = \{g \in G, \mathrm{Ad}_g(X_0) = X_0\}$ is the isotropy subgroup at X_0. Recall that U is a closed connected subgroup of the same rank as G (see e.g. [2], Chapter 8). We shall denote by $\mathfrak{u}$ the Lie algebra of U. Let T be a maximal torus in U, $\mathfrak{t}$ its Lie algebra and $\mathfrak{h} := \mathfrak{t}^{\mathbb{C}}$ the complexification of $\mathfrak{t}$, which is a Cartan subalgebra of the complex semisimple Lie algebra $\mathfrak{g}^{\mathbb{C}}$, included in $\mathfrak{u}^{\mathbb{C}}$. Let $R \subset \mathfrak{h}^* \setminus \{0\}$ be the root system of $\mathfrak{g}^{\mathbb{C}}$ with respect to $\mathfrak{h}$. For any $\alpha \in R$ denote by $\mathfrak{g}_\alpha$ the 1-dimensional root space corresponding to α, defined by

$$\mathfrak{g}_\alpha := \{X \in \mathfrak{g}^{\mathbb{C}} : \quad [H, X] = \alpha(H)X, \quad \forall H \in \mathfrak{h}\}.$$

Let

$$\begin{aligned} \mathfrak{g}^{\mathbb{C}} &= \mathfrak{h} + \sum_{\alpha \in R} \mathfrak{g}_\alpha \\ \mathfrak{u}^{\mathbb{C}} &= \mathfrak{h} + \sum_{\alpha \in R_0} \mathfrak{g}_\alpha \end{aligned}$$

be the corresponding Cartan decompositions, where $R_0 \subset R$ is the root system of $\mathfrak{u}^{\mathbb{C}}$ with respect to $\mathfrak{h}$. The roots from $R \setminus R_0$ are called complementary roots.

Since G is compact, $\mathfrak{g}$ is a compact real form of $\mathfrak{g}^{\mathbb{C}}$ and can be described in terms of a Weyl basis as follows. Recall first that choosing a Weyl basis amounts to chose root vectors $E_\alpha \in \mathfrak{g}_\alpha$ such that

$$\langle E_\alpha, E_{-\alpha} \rangle = 1, \quad \forall \alpha \in R$$

(where $\langle X, Y \rangle := \operatorname{tr}(\operatorname{ad}_X \circ \operatorname{ad}_Y)$ denotes the Killing form of $\mathfrak{g}^{\mathbb{C}}$) and

$$N_{-\alpha,-\beta} = -N_{\alpha\beta}, \quad \forall \alpha, \beta \in R \tag{3}$$

where the constants $N_{\alpha\beta}$ are defined by

$$[E_\alpha, E_\beta] = N_{\alpha\beta} E_{\alpha+\beta}, \quad \forall \alpha, \beta \in R. \tag{4}$$

With respect to a Weyl basis, $\mathfrak{g}$ is the real Lie algebra generated by $A_\alpha := E_\alpha - E_{-\alpha}$, $B_\alpha := i(E_\alpha + E_{-\alpha})$ (for any $\alpha \in R$) and $\mathfrak{t}$ (which is also the real vector space spanned by $i[E_\alpha, E_{-\alpha}]$, for all $\alpha \in R$). Similarly, $\mathfrak{u}$ is the real Lie algebra generated by A_α, B_α (for any $\alpha \in R_0$) and $\mathfrak{t}$. It follows that

$$T_e M = \mathfrak{m} := \mathfrak{g} \ominus \mathfrak{u} = \sum_{\alpha \in R \setminus R_0} \operatorname{Span}_{\mathbb{R}}\{A_\alpha, B_\alpha\} \tag{5}$$

is an Ad_T-invariant complement of $\mathfrak{u}$ in $\mathfrak{g}$, which will be identified with the tangent space T_eM at the identity coset e. By complexification,

$$T_e^{\mathbb{C}}M = \mathfrak{m}^{\mathbb{C}} = \sum_{\alpha \in R \setminus R_0} \mathfrak{g}_\alpha. \tag{6}$$

A G-invariant (or shortly, "invariant") tensor field on M is uniquely determined by its value at e, which is an Ad_T-invariant tensor on the vector space $\mathfrak{m}$.

In the proof of our main result we shall use a general formula for the exterior derivative of invariant forms on M, as follows. If $\alpha \in \Omega^k(M)$ is invariant, then $d\alpha$ is also invariant. At e,

$$(d\alpha)(X_0, \cdots, X_k) = \sum_{i<j} (-1)^{i+j} \alpha([X_i, X_j], X_1, \cdots, \widehat{X_i}, \cdots, \widehat{X_j}, \cdots, X_k), \tag{7}$$

for any $X_i \in \mathfrak{m}$. In (7) the hat means that the term is omitted; the Lie bracket $[X_i, X_j]$ of vectors $X_i, X_j \in \mathfrak{m}$ is obtained by taking the Lie bracket in $\mathfrak{g}$, followed by the standard projection $\mathfrak{g} = \mathfrak{m} \oplus \mathfrak{u} \to \mathfrak{m}$.

Invariant Hermitian and Kähler structures. Following [1], [3], [8] we develop the algebraic description of invariant Hermitian and Kähler structures on M. We preserve the notations from the previous paragraph. In addition, we fix a positive root system R_0^+ of R_0.

Let (g, J) be an invariant Hermitian structure on M. The value J_e of J at e is an Ad_T-invariant complex structure on $\mathfrak{m}$ and its complex linear extension

$$J_e^c : \mathfrak{m}^{\mathbb{C}} \to \mathfrak{m}^{\mathbb{C}}$$

is of the form

$$J_e^c E_\alpha = i\epsilon_\alpha E_\alpha, \quad \forall \alpha \in R \setminus R_0, \tag{8}$$

where $\epsilon_\alpha = \pm 1$ and $\epsilon_\alpha = -\epsilon_{-\alpha}$ for any $\alpha \in R \setminus R_0$. The set

$$\Sigma := \{\alpha \in R \setminus R_0, \quad \epsilon_\alpha = 1\}$$

determines J completely and is called the set of roots of J. Note that

$$\Sigma \cup (-\Sigma) = R \setminus R_0, \quad \Sigma \cap (-\Sigma) = \emptyset,$$

where $-\Sigma := \{-\alpha, \alpha \in \Sigma\}$. The integrability of J implies that Σ is closed and $R^+ := R_0^+ \cup \Sigma$ is a system of positive roots of R. Conversely, any closed set Σ of complementary roots such that $R_0^+ \cup \Sigma$ is a positive root system of R, defines an invariant complex structure on M (see Proposition 13.7 of [3] and [8]).

Similarly, the value g_e of the Hermitian metric g at e is a symmetric positive definite inner product $(X, Y) = -\langle \Lambda X, Y\rangle$ on $\mathfrak{m}$, where $\Lambda : \mathfrak{m} \to \mathfrak{m}$ is a linear isomorphism. Since $(\cdot, \cdot)$ is Ad_T-invariant, the complex linear extension

$$\Lambda^c : \mathfrak{m}^{\mathbb{C}} \to \mathfrak{m}^{\mathbb{C}}$$

of Λ to $\mathfrak{m}^{\mathbb{C}}$ is of the form

$$\Lambda^c(E_\alpha) = g_\alpha E_\alpha, \quad \forall \alpha \in R \setminus R_0, \tag{9}$$

for some constants g_α. Since $(\cdot, \cdot)$ is symmetric and positive definite,

$$g_\alpha = g_{-\alpha}, \quad g_\alpha > 0, \quad \forall \alpha \in R \setminus R_0. \tag{10}$$

Therefore, (g, J) is completely determined by Σ and the constants $\{g_\alpha\}_{\alpha \in R\setminus R_0}$. Conversely, any closed set $\Sigma \subset R \setminus R_0$ such that $R_0^+ \cup \Sigma$ is a positive root system of R, together with positive constants $\{g_\alpha\}_{\alpha\in R\setminus R_0}$ such that $g_\alpha = g_{-\alpha}$, for any $\alpha \in R \setminus R_0$, define an invariant Hermitian structure on M.

We now determine necessary and sufficient conditions on $\{g_\alpha\}$ such that the Hermitian pair (g, J) is Kähler. Using (8) and (9), it can be checked that the complex linear extension of the Kähler form $\omega := g(J\cdot, \cdot)$ at e, applied to a pair (E_α, E_β) of root vectors corresponding to complementary roots $\alpha, \beta \in R \setminus R_0$, is zero when $\alpha + \beta \neq 0$, and

$$\omega_e(E_\alpha, E_{-\alpha}) = (J_e E_\alpha, E_{-\alpha}) = -i g_\alpha \epsilon_\alpha, \quad \forall \alpha \in R \setminus R_0. \tag{11}$$

For any $\alpha, \beta, \gamma \in R \setminus R_0$,

$$(d\omega)(E_\alpha, E_\beta, E_\gamma) = 0 \tag{12}$$

unless $\alpha + \beta + \gamma = 0$, in which case

$$(d\omega)(E_\alpha, E_\beta, E_\gamma) = -i N_{\alpha\beta}(\epsilon_\alpha g_\alpha + \epsilon_\beta g_\beta + \epsilon_\gamma g_\gamma). \tag{13}$$

Relations (12) and (13) are a consequence of (7), (11) and the following property of $\{N_{\alpha\beta}\}$ (see Chapter 5 of [5]): if $\alpha, \beta, \gamma \in R$ are such that $\alpha + \beta + \gamma = 0$, then

$$N_{\alpha\beta} = N_{\beta\gamma} = N_{\gamma\alpha}. \tag{14}$$

Thus the pair (g, J) is Kähler if and only if

$$\epsilon_\alpha g_\alpha + \epsilon_\beta g_\beta + \epsilon_\gamma g_\gamma = 0$$

for any $\alpha, \beta, \gamma \in R \setminus R_0$ with $\alpha + \beta + \gamma = 0$; equivalently (g, J) is Kähler if and only if

$$g_{\alpha+\beta} = g_\alpha + g_\beta$$

for all $\alpha, \beta \in \Sigma$ with $\alpha + \beta \in \Sigma$. To summarize, we get the following description of invariant Kähler structures on M (for M a full flag, see [7]).

Proposition 3. *[1] Let (g, J) be an invariant Hermitian structure on M, defined by a closed set Σ and positive constants $\{g_\alpha\}_{\alpha \in R \setminus R_0}$ as above. Then (g, J) is Kähler if and only if for any $\alpha, \beta \in \Sigma$ such that $\alpha + \beta \in R$,*

$$g_{\alpha+\beta} = g_\alpha + g_\beta. \tag{15}$$

Note that for two roots $\alpha, \beta \in \Sigma$ the conditions $\alpha + \beta \in R$ and $\alpha + \beta \in \Sigma$ are equivalent, because Σ is closed.

3 Proof of the main result

In this Section we prove Theorem 1. Let $M = G/U$ be an adjoint orbit of a compact connected semisimple Lie group G and (g, J) an invariant Hermitian structure on M. We preserve the notations from the previous Section. Recall that R denotes the root system of $\mathfrak{g}^{\mathbb{C}}$ (the complexification of the Lie algebra of G) with respect to a Cartan subalgebra $\mathfrak{h}$ included in $\mathfrak{u}^{\mathbb{C}}$ (the complexification of the Lie algebra of U) and R_0^+ is a system of positive roots of the root system R_0 of $\mathfrak{u}^{\mathbb{C}}$ with respect to $\mathfrak{h}$. Thus, the pair (g, J) is defined by a closed complementary set of roots $\Sigma \subset R \setminus R_0$ and positive constants $\{g_\alpha\}_{\alpha \in R \setminus R_0}$ such that $R_0^+ \cup \Sigma := R^+$ is a positive root system of R, $g_\alpha = g_{-\alpha}$ for any $\alpha \in R \setminus R_0$ and condition (15) is satisfied.

It is easy to check, using (7), (8), (13) and (14), that for any $\alpha, \beta \in \Sigma$,

$$\begin{aligned} \frac{1}{2} dd^J\omega(E_\alpha, E_\beta, E_{-\alpha}, E_{-\beta}) = {} & N_{\alpha\beta}^2 \left(g_{\alpha+\beta} - g_\alpha - g_\beta\right) \\ & + \epsilon_{\alpha-\beta} N_{\alpha,-\beta}^2 \left(\epsilon_{\alpha-\beta} g_{\alpha-\beta} - g_\alpha + g_\beta\right), \end{aligned}$$

where $\epsilon_{\alpha-\beta} = 0$ if $\alpha - \beta \in R_0$ or if $\alpha - \beta \notin R$ and $\epsilon_{\alpha-\beta} = \pm 1$ if $\alpha - \beta \in (\pm\Sigma)$.

Assume now that $dd^J\omega = 0$. The above relation implies that

$$N^2_{\alpha\beta}\left(g_{\alpha+\beta} - g_\alpha - g_\beta\right) + \epsilon_{\alpha-\beta} N^2_{\alpha,-\beta}\left(\epsilon_{\alpha-\beta} g_{\alpha-\beta} - g_\alpha + g_\beta\right) = 0, \tag{16}$$

for any $\alpha, \beta \in \Sigma$. Using an induction argument on the length of roots of Σ with respect to the order defined by R^+ we will show that (16) implies that

$$g_{\alpha+\beta} = g_\alpha + g_\beta, \tag{17}$$

for any $\alpha, \beta \in \Sigma$ such that $\alpha + \beta \in \Sigma$. Suppose first that $\alpha, \beta \in \Sigma$ are both of length one. Then $\alpha - \beta \notin R$ and hence $N_{\alpha,-\beta} = 0$. Since $\alpha + \beta \in R$, $N_{\alpha\beta} \neq 0$ and relation (16) implies (17). Suppose now that (17) holds for all $\alpha, \beta \in \Sigma$ of length less or equal to k, such that $\alpha + \beta \in \Sigma$. Let $\gamma, \delta \in \Sigma$ be of length less or equal to $k+1$, such that $\gamma + \delta \in \Sigma$. We will show that

$$g_{\gamma+\delta} = g_\gamma + g_\delta. \tag{18}$$

If $\gamma - \delta \in R_0$ or $\alpha - \beta \notin R$, $\epsilon_{\gamma-\delta} = 0$ and from (16) again we get (18) (since $N_{\gamma\delta} \neq 0$ when $\gamma + \delta \in R$). Suppose now that $\gamma - \delta \in R \setminus R_0$. Without loss of generality, we assume moreover that $\gamma - \delta \in \Sigma$. Since $\gamma = \delta + (\gamma - \delta)$ and γ has length at most $k+1$, both δ and $\gamma - \delta$ have length less or equal to k (and belong to Σ). From our induction hypothesis,

$$g_\gamma = g_\delta + g_{\gamma-\delta}. \tag{19}$$

Using (16) for $\alpha := \gamma$ and $\beta := \delta$, together with (19), we obtain (18). Relation (17) is now proved. From Proposition 3, (g, J) is Kähler. This concludes the proof of Theorem 1.

4 Appendix: generalized Kähler geometry

Let N be a smooth manifold, $\mathbb{T}N = TN \oplus T^*N$ the generalized tangent bundle of N and g_{can} the canonical indefinite metric on $\mathbb{T}N$ given by

$$g_{\mathrm{can}}(X+\xi, Y+\eta) := \frac{1}{2}\left(\xi(Y) + \eta(X)\right), \quad \forall X+\xi, Y+\eta \in \mathbb{T}N. \tag{20}$$

Definition 4. *[6] i) An almost generalized complex structure on* N *is a* g_{can}*-skew-symmetric field of endomorphisms*

$$\mathcal{J} : \mathbb{T}N \to \mathbb{T}N$$

with $\mathcal{J}^2 = -\mathrm{Id}$ *(where "*Id*" denotes the identity endomorphism of* $\mathbb{T}N$*).*

ii) An almost generalized complex structure $\mathcal{J}$ *is integrable (or is a generalized complex structure) if the* i*-eigenbundle* $L = (\mathbb{T}N)^{1,0} \subset (\mathbb{T}N)^{\mathbb{C}}$ *of* $\mathcal{J}$ *is closed under the complex linear extension of the Courant bracket* $[\cdot, \cdot]$*, defined by*

$$[X + \xi, Y + \eta] := [X, Y] + L_X\eta - L_Y\xi - \frac{1}{2}d\left(\eta(X) - \xi(Y)\right), \tag{21}$$

for any smooth sections $X + \xi$ *and* $Y + \eta$ *of* $\mathbb{T}N$*.*

iii) A generalized Kähler structure on N *is a pair* $(\mathcal{J}_1, \mathcal{J}_2)$ *of commuting, generalized complex structures such that the bundle endomorphism* $D := -\mathcal{J}_1\mathcal{J}_2$ *of* $\mathbb{T}N$ *is positive definite with respect to the standard indefinite metric* g_{can}*, i.e. the bilinear form* $g_{\mathrm{can}} \circ D$*, defined by*

$$(g_{\mathrm{can}} \circ D)(X + \xi, Y + \eta) = g_{\mathrm{can}}(D(X + \xi), Y + \eta), \quad \forall X + \xi, Y + \eta \in \mathbb{T}N$$

is positive definite.

Generalized geometry and bi-Hermitian geometry are related as follows.

Theorem 5. *[4] On a manifold* N *there is a one to one correspondence between generalized Kähler structures* $(\mathcal{J}_1, \mathcal{J}_2)$ *and bi-Hermitian structures* (g, J_+, J_-) *together with a 2-form* $b \in \Omega^2(N)$*, such that the condition*

$$db = -d^{J_+}\omega_+ = d^{J_-}\omega_- \tag{22}$$

holds. Here $\omega_\pm := g(J_\pm \cdot, \cdot)$ *are the Kähler forms of* $(J_\pm, g)$*.*

Proof. We briefly explain how $(\mathcal{J}_1, \mathcal{J}_2)$ determines (g, J_+, J_-, b). Define $D := -\mathcal{J}_1\mathcal{J}_2$. Since $\mathcal{J}_1$ and $\mathcal{J}_2$ commute, $D^2 = \mathrm{Id}$. Let $C_\pm$ be the eigenbundles of D, with eigenvalues ± 1. They are preserved by both $\mathcal{J}_1$ and $\mathcal{J}_2$ and the standard projections

$$\pi_\pm : C_\pm \to TN$$

are isomorphisms. The almost complex structures $J_\pm$ of N are induced, by means of $\pi_\pm$, by the restriction of $\mathcal{J}_1$ to $C_\pm$. The bi-Hermitian metric and the b-field can also be read from $C_\pm$: the eigenbundle $C_\pm$ is the graph of $b \pm g : TN \to T^*N$ (here b associates to $X \in TN$ the 1-form $i_X b := b(X, \cdot)$ and similarly for g). According to [4], g is a Riemannian metric compatible with $J_{\pm}$ (i.e. $J_{\pm}$ are g-skew-symmetric) and the integrability of $\mathcal{J}_1$ and $\mathcal{J}_2$ is equivalent to the integrability of the almost complex structures $J_{\pm}$, together with condition (22).

$\square$

References

[1] D. V. Alekseevsky, A. M. Perelomov: *Invariant Kähler-Einstein metrics on compact homogeneous spaces*, Funct. Anal. Applic. 20 (1986), no. 3, p. 171-182.

[2] A. Besse: *Einstein manifolds*, Springer-Verlag, 1987.

[3] A. Borel, F. Hirzebruch: *Characteristic classes and homogeneous spaces I*, American J. Math., vol. 80 (1958), p. 459-536.

[4] M. Gualtieri: *Generalized Complex Geometry*, D.Phil thesis 2003, University of Oxford, UK.

[5] S. Helgason: ”Differential Geometry, Lie groups and Symmetric Spaces”, Academic Press, New York, 1978.

[6] N. Hitchin: *Generalized Calabi-Yau manifolds*, Quart. J. Math., 54 no. 3 (2003), p. 281-308.

[7] L. A. B. San Martin, C. J.C. Negreiros: *Invariant almost Hermitian structures on flag manifolds*, Adv. Math. 178 (2003), p. 277-310.

[8] H. C. Wang: *Closed manifolds with homogeneous complex structures*, Amer. J. Math. 76 (1954), p. 1-32.

DMITRI V. ALEKSEEVSKY: Edinburgh University and Maxwell Institute of Mathematical Study, The King’s Buildings, JCMB, Mayfield Road, Edinburgh, EH9 3JZ,UK, D.Aleksee@ed.ac.uk

LIANA DAVID: Institute of Mathematics ”Simion Stoilow” of the Romanian Academy; Calea Grivitei nr. 21, Sector 1, Bucharest, Romania; liana.david@imar.ro